

Table 5.12a
Adjusted Patient Survival, Deceased Donor non-ECD Kidney Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2006 - 2007)			(Tx 2006 - 2007)			(Tx 2002 - 2007)			(Tx 1997 - 2007)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
Total												
All	14,409	98.7%	0.1%	14,409	96.9%	0.2%	39,452	85.8%	0.3%	68,081	64.8%	0.3%
Age at Tx												
<1 Year	2	+	+	2	+	+	4	+	+	5	+	+
1-5 Years	152	98.7%	0.9%	152	97.5%	1.3%	343	94.0%	1.6%	533	89.4%	1.9%
6-11 Years	198	99.5%	0.5%	198	99.0%	0.7%	521	96.7%	0.9%	829	93.9%	1.4%
12-17 Years	583	99.7%	0.2%	583	98.8%	0.5%	1,368	94.3%	1.0%	2,053	84.9%	1.5%
18-34 Years	1,675	99.2%	0.2%	1,675	98.7%	0.3%	5,058	92.5%	0.5%	9,722	81.8%	0.6%
35-49 Years	4,077	99.1%	0.1%	4,077	98.0%	0.2%	11,602	89.2%	0.4%	21,370	71.6%	0.5%
50-64 Years	5,670	98.1%	0.2%	5,670	95.7%	0.3%	15,504	82.5%	0.4%	25,900	55.5%	0.6%
65+ Years	2,052	97.5%	0.3%	2,052	93.3%	0.6%	5,052	69.6%	1.0%	7,669	34.8%	1.2%
Recipient Ethnicity/Race												
White	6,360	98.8%	0.1%	6,360	96.8%	0.2%	18,165	84.8%	0.4%	33,307	63.8%	0.5%
African American	4,631	98.4%	0.2%	4,631	96.6%	0.3%	12,442	84.1%	0.5%	20,847	61.0%	0.6%
Hispanic/Latino	2,320	98.7%	0.2%	2,320	97.4%	0.3%	5,924	90.2%	0.5%	9,220	72.2%	0.9%
Asian	878	98.9%	0.3%	878	97.7%	0.5%	2,377	90.3%	0.8%	3,867	74.8%	1.3%
Other/Multi-race	220	99.3%	0.5%	220	98.2%	0.8%	543	88.3%	1.8%	837	70.9%	2.8%
Unknown	0	-	-	0	-	-	1	+	+	3	+	+
Recipient Gender												
Female	5,624	98.8%	0.1%	5,624	97.3%	0.2%	15,673	87.0%	0.4%	27,089	66.4%	0.5%
Male	8,785	98.6%	0.1%	8,785	96.7%	0.2%	23,779	85.0%	0.3%	40,992	63.8%	0.4%

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether kidneys were allocated under ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

Table 5.12a (Continued)
Adjusted Patient Survival, Deceased Donor non-ECD Kidney Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2006 - 2007)			(Tx 2006 - 2007)			(Tx 2002 - 2007)			(Tx 1997 - 2007)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
Total												
All	14,409	98.7%	0.1%	14,409	96.9%	0.2%	39,452	85.8%	0.3%	68,081	64.8%	0.3%
Primary Diagnosis												
Glomerular Diseases	3,322	99.0%	0.2%	3,322	97.8%	0.3%	9,466	88.6%	0.5%	17,073	72.0%	0.6%
Diabetes	3,422	98.1%	0.2%	3,422	95.3%	0.4%	9,371	78.9%	0.6%	15,697	49.2%	0.7%
Hypertensive												
Nephrosclerosis	3,591	98.5%	0.2%	3,591	96.6%	0.3%	8,914	84.6%	0.6%	14,194	62.0%	0.8%
Polycystic Kidneys	1,256	99.3%	0.2%	1,256	98.0%	0.4%	3,495	92.9%	0.6%	6,336	78.3%	0.8%
Tubular and Interstitial Diseases	649	99.0%	0.4%	649	97.6%	0.6%	1,932	88.2%	0.9%	3,600	69.7%	1.3%
Renovascular & Other Vascular Diseases	487	97.6%	0.6%	487	95.3%	0.9%	1,848	84.9%	1.0%	3,488	61.5%	1.4%
Congenital, Rare Familial, & Metabolic Disorders	427	98.8%	0.6%	427	97.8%	0.9%	1,101	91.2%	1.3%	1,819	74.6%	2.2%
Neoplasms	61	97.5%	1.8%	61	97.5%	1.8%	136	81.4%	5.2%	212	51.6%	6.5%
Other	1,187	99.0%	0.3%	1,187	98.0%	0.4%	3,171	87.0%	0.8%	5,633	69.6%	1.0%
Unknown	7	81.2%	16.0%	7	81.2%	16.0%	18	44.4%	18.2%	29	69.1%	9.8%

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether kidneys were allocated under ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

Table 5.12a (Continued)
Adjusted Patient Survival, Deceased Donor non-ECD Kidney Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2006 - 2007)			(Tx 2006 - 2007)			(Tx 2002 - 2007)			(Tx 1997 - 2007)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
Total												
All	14,409	98.7%	0.1%	14,409	96.9%	0.2%	39,452	85.8%	0.3%	68,081	64.8%	0.3%
Transplant Center State												
Alabama	249	98.9%	0.6%	249	97.4%	1.0%	650	85.7%	1.8%	1,287	64.5%	2.1%
Arkansas	157	99.4%	0.6%	157	97.8%	1.1%	395	82.7%	3.2%	591	57.2%	3.8%
Arizona	234	100.0%	0.0%	234	97.7%	0.9%	595	92.5%	1.3%	978	77.4%	2.1%
California	1,768	99.0%	0.2%	1,768	97.6%	0.3%	4,828	87.3%	0.7%	8,025	68.5%	0.9%
Colorado	163	99.5%	0.5%	163	97.1%	1.2%	530	89.0%	1.7%	879	65.1%	2.8%
Connecticut	95	99.1%	0.8%	95	97.4%	1.5%	329	87.8%	2.3%	549	68.9%	3.3%
District of Columbia	135	98.7%	0.9%	135	96.7%	1.5%	404	84.2%	2.8%	687	73.6%	2.5%
Delaware	9	+	+	9	+	+	14	+	+	23	+	+
Florida	949	98.7%	0.3%	949	97.5%	0.5%	2,634	85.9%	1.0%	4,616	65.3%	1.2%
Georgia	394	98.9%	0.5%	394	96.8%	0.8%	1,064	87.0%	1.4%	1,931	64.7%	1.9%
Hawaii	38	100.0%	0.0%	38	98.2%	1.8%	144	91.8%	2.8%	243	71.9%	6.1%
Iowa	111	98.9%	0.8%	111	98.3%	1.0%	330	83.7%	2.5%	580	62.5%	3.5%
Illinois	446	99.2%	0.4%	446	98.0%	0.6%	1,387	86.7%	1.2%	2,473	63.2%	1.8%
Indiana	179	98.8%	0.7%	179	97.1%	1.1%	713	90.6%	1.3%	1,185	76.1%	2.1%
Kansas	115	99.3%	0.7%	115	98.6%	1.0%	336	91.9%	1.9%	569	68.5%	3.5%
Kentucky	133	98.8%	0.8%	133	97.6%	1.2%	446	84.2%	2.4%	847	53.8%	3.0%
Louisiana	268	99.0%	0.5%	268	97.1%	0.9%	858	82.2%	1.9%	1,419	59.3%	2.5%
Massachusetts	273	97.2%	0.9%	273	94.7%	1.3%	770	89.6%	1.3%	1,434	64.1%	2.1%
Maryland	307	98.1%	0.7%	307	95.0%	1.1%	793	80.6%	1.8%	1,562	55.8%	2.3%
Maine	59	100.0%	0.0%	59	97.3%	1.9%	158	90.6%	2.5%	270	72.4%	4.6%
Michigan	484	97.9%	0.6%	484	95.7%	0.8%	1,326	84.0%	1.3%	2,239	61.7%	1.8%
Minnesota	221	99.3%	0.5%	221	97.6%	0.9%	578	88.8%	1.5%	1,072	67.9%	2.3%
Missouri	432	99.6%	0.3%	432	98.0%	0.6%	1,046	88.7%	1.4%	1,885	66.3%	1.9%
Mississippi	100	98.9%	1.1%	100	97.8%	1.5%	195	86.9%	3.5%	313	67.0%	5.2%
North Carolina	469	97.7%	0.6%	469	96.6%	0.8%	1,119	88.9%	1.3%	1,787	67.9%	2.1%
North Dakota	34	98.0%	2.0%	34	95.9%	2.8%	105	80.0%	4.3%	168	58.9%	6.5%
Nebraska	82	99.1%	0.9%	82	98.1%	1.3%	200	86.9%	2.9%	369	69.3%	4.2%
New Hampshire	26	+	+	26	+	+	87	88.4%	3.8%	136	80.9%	4.2%
New Jersey	301	95.8%	1.0%	301	93.4%	1.3%	784	80.0%	2.0%	1,384	60.6%	2.3%
New Mexico	85	+	+	85	+	+	197	98.0%	1.0%	342	69.1%	4.4%
Nevada	93	97.2%	1.6%	93	93.5%	2.4%	250	77.5%	3.6%	416	57.6%	4.1%

(Continued)

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether kidneys were allocated under ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

Table 5.12a (Continued)
Adjusted Patient Survival, Deceased Donor non-ECD Kidney Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2006 - 2007)			(Tx 2006 - 2007)			(Tx 2002 - 2007)			(Tx 1997 - 2007)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
Transplant Center State												
New York	954	98.2%	0.4%	954	95.8%	0.6%	2,191	83.3%	1.2%	3,712	65.8%	1.3%
Ohio	487	98.2%	0.5%	487	96.5%	0.8%	1,340	82.0%	1.4%	2,651	59.2%	1.6%
Oklahoma	127	99.3%	0.7%	127	97.1%	1.4%	426	83.8%	2.2%	761	61.6%	2.9%
Oregon	177	99.2%	0.6%	177	98.3%	0.8%	485	84.6%	2.2%	820	66.0%	3.0%
Pennsylvania	993	99.0%	0.3%	993	96.7%	0.5%	2,780	83.7%	0.9%	4,736	63.8%	1.2%
Puerto Rico	120	99.1%	0.9%	120	98.2%	1.3%	324	92.6%	1.8%	473	56.9%	6.1%
Rhode Island	42	100.0%	0.0%	42	98.1%	1.9%	148	84.0%	3.2%	251	58.2%	5.8%
South Carolina	229	100.0%	0.0%	229	95.5%	1.3%	559	86.6%	2.1%	986	57.9%	3.0%
South Dakota	38	100.0%	0.0%	38	98.4%	1.6%	97	90.3%	3.5%	179	74.8%	4.3%
Tennessee	414	98.0%	0.6%	414	95.9%	0.9%	1,119	83.7%	1.4%	1,840	55.9%	2.3%
Texas	1,224	98.1%	0.4%	1,224	96.9%	0.5%	3,423	85.7%	0.8%	5,860	66.7%	1.1%
Utah	107	100.0%	0.0%	107	99.3%	0.7%	310	93.3%	1.7%	551	76.8%	2.9%
Virginia	342	98.0%	0.7%	342	96.7%	0.9%	846	83.9%	1.7%	1,469	60.1%	2.2%
Vermont	40	100.0%	0.0%	40	94.9%	2.9%	75	90.9%	3.2%	114	85.4%	4.4%
Washington	280	99.7%	0.3%	280	98.9%	0.6%	892	88.7%	1.3%	1,423	64.1%	2.6%
Wisconsin	354	99.3%	0.4%	354	97.5%	0.8%	976	84.9%	1.5%	1,618	65.4%	2.1%
West Virginia	72	98.9%	1.1%	72	94.3%	2.5%	196	77.3%	4.0%	378	54.4%	4.2%

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether kidneys were allocated under ECD system. Non-ECDs include all other donors, including SCDs and DCDs.